

---

We never meant to design Field Gothic. All we wanted was a version of our Pressio family with conventional rounded counters, which we thought might sell a bit better to the corner-averse. How long could that take, a couple weeks? And then we realized that, without those squared-off counters, we could do extra-light and hairline weights. And then we began to wonder what wider styles might look like. And then the WIP began to remind us of things: Unimark subway signage, midcentury Geigy promotions, old Westinghouse lightbulb packs, our dad's Columbia and Blue Note jazz albums, Letraset Compacta, the Schelter & Giesecke specimens we studied when we started getting serious about drawing type. And then we were off to the races. At the time of writing, Field Gothic has snowballed into an eight-weight, eight-width, possibly-too-personal journey through the history of American and European sanses that may or may not parallel our own journey from New York City to Dublin. Overall, the look is crisp and on the cool side, with slightly superelliptical curves. The lightest weights are almost monoline. Above that, arches and bowls have sharply contrasting junctures and a notably snug fit. At 128 styles (including matching italics), it's by far the largest family we've ever designed, or ever want to design. We are very, very tired.

---

Field Gothic N010 & Italic

Field Gothic N011 & Italic

Field Gothic N012 & Italic

Field Gothic N013 & Italic

Field Gothic N014 & Italic

Field Gothic N015 & Italic

Field Gothic N016 & Italic

Field Gothic N017 & Italic

Field Gothic N020 & Italic

Field Gothic N021 & Italic

Field Gothic N022 & Italic

Field Gothic N023 & Italic

Field Gothic N024 & Italic

Field Gothic N025 & Italic

Field Gothic N026 & Italic

Field Gothic N027 & Italic

Field Gothic N030 & Italic

Field Gothic N031 & Italic

Field Gothic N032 & Italic

Field Gothic N033 & Italic

Field Gothic N034 & Italic

Field Gothic N035 & Italic

Field Gothic N036 & Italic

Field Gothic N037 & Italic

Field Gothic N040 & Italic

Field Gothic N041 & Italic

Field Gothic N042 & Italic

Field Gothic N043 & Italic

Field Gothic N044 & Italic

Field Gothic N045 & Italic

Field Gothic N046 & Italic

Field Gothic N047 & Italic

Field Gothic N050 & Italic

Field Gothic N051 & Italic

Field Gothic N052 & Italic

Field Gothic N053 & Italic

**Field Gothic N054 & Italic**

**Field Gothic N055 & Italic**

**Field Gothic N056 & Italic**

**Field Gothic N057 & Italic**

Field Gothic N060 & Italic

Field Gothic N061 & Italic

Field Gothic N062 & Italic

Field Gothic N063 & Italic

**Field Gothic N064 & Italic**

**Field Gothic N065 & Italic**

**Field Gothic N066 & Italic**

**Field Gothic N067 & Italic**

Field Gothic N070 & Italic

Field Gothic N071 & Italic

Field Gothic N072 & Italic

Field Gothic N073 & Italic

**Field Gothic N074 & Italic**

**Field Gothic N075 & Italic**

**Field Gothic N076 & Italic**

**Field Gothic N077 & Italic**

Field Gothic N080 & Italic

Field Gothic N081 & Italic

Field Gothic N082 & Italic

**Field Gothic N083 & Italic**

**Field Gothic N084 & Italic**

**Field Gothic N085 & Italic**

**Field Gothic N086 & Italic**

**Field Gothic N087 & Italic**

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*



---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other,

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other,*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*



---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the*

36pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condltion. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other,*

48pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the*

36pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

48pt

---

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*



---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked*

---

36pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the

---

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked*

---

36pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and***

---

48pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which***



---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may

---

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured

---

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured

---

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has*



---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are

---

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which*

---

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromato-

---

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

---

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromato-*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which*

---

24pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-**

---

34pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up**

---

48pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which**

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which*



---

24pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross**

---

34pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,**

---

48pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,**

---

24pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross*

---

34pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,*

---

48pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,*

---

24pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross**

---

34pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,**

---

48pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,**

---

24pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross***

---

34pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,***

---

48pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,***

---

21pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,

---

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and general-

---

21pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,*

---

48pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and general-*



---

21pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,

---

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

---

21pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,*

---

48pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-*

---

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

---

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a sub-

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

---

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

---

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a*

18pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,*

---

24pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a sub-*

28pt

*THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless*

---

48pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-*

---

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

---

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

---

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

---

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a*

18pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,*

---

24pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a sub-*

28pt

*THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless*

---

48pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-*



14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gen-

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a*

18pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-*

24pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance***

28pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless***

48pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gen-***

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,*

18pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal*

24pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance***

28pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless***

48pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and***

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,**

18pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal**

24pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance**

28pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colour-**

48pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and**

14pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,***

18pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal***

24pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance***

28pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colour-***

48pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and***



14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies**

18pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal**

24pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of**

28pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a col-**

48pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and**

14pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies***

18pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal***

24pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of***

28pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a col-***

48pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and***

---

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is

---

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,

---

18pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is*

---

48pt

*THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,*

---

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it,

---

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,

---

18pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it,*

---

48pt

*THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,*



9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an eye-spot,

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral,

9pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an eye-spot,*

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes*

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is*

18pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily*

48pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral,***

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

48pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral**

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

9pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an eye-spot,*

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely*

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes*

18pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily*

48pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral***

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral



9pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the*

14pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely***

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes*

18pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily***

48pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral***



9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral

9pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the*

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which some-*

14pt

***THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely***

18pt

***THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily***

48pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral***

9pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these**

18pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily**

48pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiven-**

9pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the*

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,*

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these*

18pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily*

48pt

***THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiven-***

9pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these**

18pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,**

48pt

**THE INDIVIDUALS ARE OF A VERY variable form, but are always unicellular, and floating about in free condition. Cell is dorsiven-**



9pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially**

18pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,**

48pt

**THE INDIVIDUALS ARE OF A VERY variable form, but are always unicellular, and floating about in free condition. Cell is dorsiven-**



---

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no silica

---

46pt

THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is dorsiven-

---

16pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no silica*

---

46pt

*THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is dorsiven-*

---

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no

---

46pt

THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is

---

16pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow*

---

32pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no*

---

46pt

*THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is*

---

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the prod-

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

---

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

---

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregular-

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.



---

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

---

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregular-

---

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.

---

9pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation*

11pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a*

---

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia*

16pt

*THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly*

---

46pt

***THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.***

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregular pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow,**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The**

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregular pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow,**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The**

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The**



9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The**



9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.**

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.**

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.**

9pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the**

11pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

16pt

**THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly**

46pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.**

---

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and generally surrounded by a colourless mem-

---

46pt

THE INDIVIDUALS ARE OF variable form, but always unicellular, and floating in a free condition. The cell is

---

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches

---

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and generally surrounded by a colourless mem-

---

46pt

THE INDIVIDUALS ARE OF variable form, but always unicellular, and floating in a free condition. The cell is



---

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow).

---

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica embedded

---

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow).

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica em-

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.

8pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and*

10pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow*

14pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-*

19pt

*THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no*

46pt

**THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.**

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-

8pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil*

10pt

*THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in*

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-**

19pt

**THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no**

46pt

**THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-**



8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-

8pt

*THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phyco-pyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless*

10pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks**

19pt

**THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which**

46pt

**THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-**

8pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

10pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks**

19pt

**THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which**

46pt

**THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free con-**

8pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-**

10pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to**

14pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks**

19pt

**THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which**

46pt

**THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free con-**

10pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-**

12pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one**

15pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is provided with**

20pt

**THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane,**

46pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free**

10pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may**

12pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one**

15pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is provided with**

20pt

**THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane,**

46pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free**



10pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,**

12pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft**

15pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is pro-**

20pt

**THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colour-**

46pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free**

10pt

**THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,**

12pt

**THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft**

15pt

**THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is pro-**

20pt

**THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colour-**

46pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free**

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longi-tudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of

---

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and gener-

---

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free condi-

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longi-tudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of

---

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and gener-

---

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free condi-

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the mem-

---

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dor-siventral, asymmetric and gen-

---

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free con-

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the mem-

---

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dor-siventral, asymmetric and gen-

---

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free con-



---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a

---

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and

---

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free

---

16pt

*THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft*

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dor-siventral, asymmetric and gen-**

---

46pt

**INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free con-**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and**

---

46pt

**INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free**

---

16pt

*THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a*

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and**

---

46pt

**INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and**

---

46pt

**INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and**

---

46pt

**INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free**



---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each**

---

32pt

**THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-**

---

46pt

**INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a**



---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed

---

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiventral, bilat-

---

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed

---

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiventral, bilat-

---

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

---

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-

---

46pt

THE INDIVIDUALS are of very variable form, but floating in a free condition.

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

---

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiventral,

---

46pt

THE INDIVIDUALS are of very variable form, but floating in a free condition.

---

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

---

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiven-

---

46pt

THE MEMBRANE will easily break up into pieces. In the forms which have

---

16pt

*THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two*

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiven-**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which have**



---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiven-**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which have**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

16pt

---

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**



---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance similar to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

16pt

**THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance similar to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which**

---

32pt

**THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell**

---

46pt

**THE MEMBRANE will easily break up into pieces. In the forms which**

---

Tabular Figures      1234567890 → 1234567890

---

Arbitrary Fractions      1/9 3/8 25/32 → 1<sub>9</sub> 3<sub>8</sub> 25<sub>32</sub>

---

Superiors & Inferiors      Estuarial<sup>3</sup> H<sub>2</sub>O → Estuarial<sup>3</sup> H<sub>2</sub>O

---

Ordinals      23<sup>a</sup> 65<sup>o</sup> → 23<sup>a</sup> 65<sup>o</sup>

---

Case-sensitive Forms      (NON-COM) «OBOE» → (NON-COM) «OBOE»

---

Ligatures      Define flee office afflict ruff Kafka surfboard fjord hofhaus  
→ Define flee office afflict ruff Kafka surfboard fjord hofhaus

---

Stylistic Set 1      Great days → Great days

---

Stylistic Set 2      Cracked jars → Cracked jars

---

Stylistic Set 3      Jolly FLAPJACK → Jolly FLAPJACK

---

Stylistic Set 4      €23.75 → €23.75





---

Designed by Max Phillips.

Design support and font engineering  
by Sebastian Carewe.

Not all applications support OpenType  
features, and not all OpenType-aware  
applications support all OpenType features.  
OpenType is either a registered trademark  
or trademark of Microsoft Corporation  
in the United States and/or other countries.

Text adapted from  
*A Handbook of Systematic Botany*  
by Dr. Johannes Eugenius Warming  
Translated from the Danish  
by Michael Cresse Potter, M.A. F.L.S.  
Published by Swan Sonnenschein in 1895

©2022–2025 Signal Type Foundry Limited  
Dublin, Ireland  
Field Gothic is a trademark  
of Signal Type Foundry Limited.  
All rights reserved.

[hello@signalfoundry.com](mailto:hello@signalfoundry.com)