
We never meant to design Field Gothic. All we wanted was a version of our Pressio family with conventional rounded counters, which we thought might sell a bit better to the corner-averse. How long could that take, a couple weeks? And then we realized that, without those squared-off counters, we could do extra-light and hairline weights. And then we began to wonder what wider styles might look like. And then the WIP began to remind us of things: Unimark subway signage, midcentury Geigy promotions, old Westinghouse lightbulb packs, our dad's Columbia and Blue Note jazz albums, Letraset Compacta, the Schelter & Giesecke specimens we studied when we started getting serious about drawing type. And then we were off to the races. A year and a half later, Field Gothic has snowballed into an eight-weight, eight-width, possibly-too-personal journey through the history of American and European sanses that may or may not parallel our own journey from New York City to Dublin. Overall, the look is crisp and on the cool side, with slightly superelliptical curves. The lightest weights are almost monoline. Above that, arches and bowls have sharply contrasting junctures and a notably snug fit. At 64 styles (128, if we ever get round to making matching italics), it will be the largest family we've ever designed, or ever want to design. We are very, very tired.

Field Gothic N010

Field Gothic N011

Field Gothic N012

Field Gothic N013

Field Gothic N014

Field Gothic N015

Field Gothic N016

Field Gothic N017

Field Gothic N020

Field Gothic N021

Field Gothic N022

Field Gothic N023

Field Gothic N024

Field Gothic N025

Field Gothic N026

Field Gothic N027

Field Gothic N030

Field Gothic N031

Field Gothic N032

Field Gothic N033

Field Gothic N034

Field Gothic N035

Field Gothic N036

Field Gothic N037

Field Gothic N040

Field Gothic N041

Field Gothic N042

Field Gothic N043

Field Gothic N044

Field Gothic N045

Field Gothic N046

Field Gothic N047

Field Gothic N050

Field Gothic N051

Field Gothic N052

Field Gothic N053

Field Gothic N054

Field Gothic N055

Field Gothic N056

Field Gothic N057

Field Gothic N060

Field Gothic N061

Field Gothic N062

Field Gothic N063

Field Gothic N064

Field Gothic N065

Field Gothic N066

Field Gothic N067

Field Gothic N070

Field Gothic N071

Field Gothic N072

Field Gothic N073

Field Gothic N074

Field Gothic N075

Field Gothic N076

Field Gothic N077

Field Gothic N080

Field Gothic N081

Field Gothic N082

Field Gothic N083

Field Gothic N084

Field Gothic N085

Field Gothic N086

Field Gothic N087

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other,

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (*Exuviella*), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which exter-

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. THE CELL IS DORSIVENTRAL, BILATERAL, ASYMMETRIC and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the

36pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromato-

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING ABOUT IN FREE CONDITION. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross

34pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,

48pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane,

21pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and general-

21pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane,

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a sub-

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gener-

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and gen-

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal

24pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colour-

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies

18pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal

24pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of

28pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a col-

48pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it,

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral,

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND FLOATING about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an eye-spot,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral,

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one; in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the cilia-cleft. In some an

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiven-

9pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and starch. Cell-nucleus one: in Polydinida several nuclei are found; contractile vacuoles many, which partly open in the

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these

18pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders,

48pt

THE INDIVIDUALS ARE OF A VERY variable form, but are always unicellular, and floating about in free condition. Cell is dorsiven-

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross fur-

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no silica

46pt

THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is dorsiven-

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, AND floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and surrounded by a colourless membrane, which has no

46pt

THE INDIVIDUALS ARE OF A variable form, but always unicellular, and floating in a free condition. The cell is

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyl, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the prod-

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches cross-wise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating in a free condition.

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregular pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow,

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition. The

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.

9pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which sometimes may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-

11pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly

46pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular and floating in a free condition.

16pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches

32pt

THE INDIVIDUALS ARE OF VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, asymmetric and generally surrounded by a colourless mem-

46pt

THE INDIVIDUALS ARE OF variable form, but always unicellular, and floating in a free condition. The cell is

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow).

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregularly-shaped

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no silica embedded

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil and

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which has no

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condition.

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless oil

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks into irregu-

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-

8pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimilation which consist of yellow, red or colourless

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell,

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free condi-

8pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT ALWAYS UNICELLULAR, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-shaped chromatophore. The coloring material (pyrophyll) consists, in addition to a modification of chlorophyll, also of phycopyrrin and peridinin; this colour is sometimes more or less masked by the products of assimila-

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the

14pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks

19pt

THE INDIVIDUALS ARE OF A very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colourless membrane, which

46pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular and floating in free con-

10pt

THE INDIVIDUALS ARE OF A VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs, which may coalesce and become a star-

12pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one

15pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is provided with

20pt

THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane,

46pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free

10pt

THE INDIVIDUALS ARE OF VERY VARIABLE FORM, BUT always unicellular, and floating about in free condition. The cell is generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of these cilia projects freely and is directed longitudinally to the front or to the rear, the other one stretches crosswise and lies close to the cell, often in a furrow (cross furrow). The chromatophores are coloured brown or green and may either be two parallel (Exuviella), or several radially placed, discs,

12pt

THE INDIVIDUALS ARE OF A VERY VARIABLE form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft

15pt

THE INDIVIDUALS ARE OF A VERY variable form, but always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but instead is formed of a substance allied to cellulose. The membrane, which externally is pro-

20pt

THE INDIVIDUALS ARE OF a very variable form, but they are always unicellular, and floating about in free condition. The cell is dorsiventral, bilateral, asymmetric and usually surrounded by a colour-

46pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular and floating in free

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longi-tudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the membrane; one of

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and gener-

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free condi-

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dor-siventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a cleft in the mem-

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dor-siventral, asymmetric and gen-

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free con-

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project through a

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project

32pt

THE INDIVIDUALS ARE OF A very variable form, but always unicellular, and floating about in a free condition. The cell is dorsiventral, asymmetric and

46pt

INDIVIDUALS ARE OF a variable form, but always unicellular, and floating in a free

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each

32pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each other, and project

32pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT ALWAYS unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed where these cross each

32pt

THE INDIVIDUALS ARE OF A very variable form, but are always unicellular, and floating about in a free condition. The cell is dorsiventral, bilat-

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two cilia are fixed

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiventral, bilat-

46pt

INDIVIDUALS ARE of a variable form, but always unicellular, and floating in a

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

32pt

THE INDIVIDUALS ARE OF a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-

46pt

THE INDIVIDUALS are of very variable form, but floating in a free condition.

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows, two

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dorsiven-

46pt

THE MEMBRANE will easily break up into pieces. In the forms which have

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross furrows,

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-

46pt

THE MEMBRANE will easily break up into pieces. In the forms which

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and cross fur-

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is dor-

46pt

THE MEMBRANE will easily break up into pieces. In the forms which

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which have longitudinal and

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell is

46pt

THE MEMBRANE will easily break up into pieces. In the forms which

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance allied to cellulose. The membrane, which externally is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell

46pt

THE MEMBRANE will easily break up into pieces. In the forms which

16pt

THE INDIVIDUALS ARE OF A VARIABLE FORM, BUT always unicellular, and floating about in a free condition. The cell is dorsiventral, bilateral, asymmetric and generally surrounded by a colourless membrane, which has no silica embedded in it, but is formed of a substance similar to cellulose. The membrane, which is provided with pores and raised borders, easily breaks up into irregularly-shaped pieces. In the forms which

32pt

THE INDIVIDUALS ARE of a very variable form, but are always unicellular, and floating about in free condition. The cell

46pt

THE MEMBRANE will easily break up into pieces. In the forms which

Tabular Figures 1234567890 → 1234567890

Arbitrary Fractions 1/9 3/8 25/32 → 1₉ 3₈ 25₃₂

Superiors & Inferiors Estuarial³ H₂O → Estuarial³ H₂O

Ordinals 23^a 65^o → 23^a 65^o

Case-sensitive Forms (NON-COM) «OBOE» → (NON-COM) «OBOE»

Ligatures Define flee office afflict ruff Kafka surfboard fjord hofhaus
→ Define flee office afflict ruff Kafka surfboard fjord hofhaus

Stylistic Set 1 Great days → Great days

Stylistic Set 2 Cracked jars → Cracked jars

Stylistic Set 3 Jolly FLAPJACK → Jolly FLAPJACK

Stylistic Set 4 €23.75 → €23.75

Designed by Max Phillips.

Not all applications support OpenType features, and not all OpenType-aware applications support all OpenType features. OpenType is either a registered trademark or trademark of Microsoft Corporation in the United States and/or other countries.

Text adapted from *A Handbook of Systematic Botany*
by Dr. Johannes Eugenius Warming
Translated from the Danish by Michael Cresse Potter, M.A. F.L.S.
Published by Swan Sonnenschein in 1895

©2022 Signal Type Foundry Limited
Dublin, Ireland
Field Gothic is a trademark
of Signal Type Foundry Limited.
All rights reserved.

hello@signalfoundry.com